



Research Article

ON INVARIANT LIGHT

Sergey Orlov¹

¹Petrozavodsk State University

Correspondence should be addressed to **Sergey Orlov**

Received March 09, 2015; Accepted March 21, 2015; Published March 23, 2015;

Copyright: © 2015 **Sergey Orlov** et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite This Article: Orlov, S. (2015). On invariant light. The International Journal of Physics, 1(1). 1-4

ABSTRACT

A new physical model of the appearance of light, properties, and its trajectory. Explains the paradox of invariance and isotropy of light based on the laws of classical physics.

KEY WORDS: Theory of vortex gravitation, cosmology, cosmogony. Aerodynamics. Invariance of light.

INTRODUCTION

For many years, scientists - physicists have tried to measure the speed of light. Galileo - in the seventeenth century. Early experiment to measure the speed of light was held Ole Roemer, a Danish physicist, in 1676. Another, more accurate way to measure the speed of light is made in Europe Hippolyte Fizeau in 1849. Francois Arago determines the speed of light in 1838 by a rotating mirror. In 1862, on the production of the speed of light worked Leon Foucault. In 1926, a result of years of efforts Albert Michelson [1] has had some very high accuracy of the speed of light:

$$c = 299\,796\,000 \pm 4\,000 \text{ m/s}$$

The main and paradoxical conclusion in the study of light was his invariance. In other words, the speed of light is the same in all inertial frames of reference is independent of the velocity of the observer and emitter. This surprising fact for the classical physics of his time for the first time proved the Michelson experiment: the independence of the speed of light on the direction (isotropic) and the orbital motion of the Earth around the sun. In the future, this paradox was confirmed by astronomers. In particular, Willem de Sitter in monitoring the spectral binaries found that the speed of light flux from removing the stars and approaches are constant speed of light (c) and equal to each

other. That is, they do not depend on the rate of the star (light sources). [2] From the point of view of classical physics this paradox so far can't be explained. Therefore, on the basis of the unknown properties of light in 1905, Einstein proposed the special theory of relativity (STR) in the paper "On the Electrodynamics of Moving Bodies" [3]. SRT conclusions were as paradoxical as the invariance of light. On the basis of the SRT can assert that the events simultaneous in one inertial reference frame will not be simultaneous in another, if these frames of reference are moving relative to each other.

In this article we propose to explain the origin, properties and the invariance of light, in full compliance with the classical laws of physics using the theory of vortex gravitation, cosmology and cosmogony. [4]

LUMINIFEROUS ETHER


In the theory of vortex gravitation, cosmology and cosmogony all celestial bodies (matter) are essential vortices. The values of the bodies (system of bodies) and the corresponding vortices may vary to an infinite value. The biggest ethereal vortex that people can watch this universal whirlwind, the smallest - atomic.

The orbital speed of ether in each vortex increases towards the center of an inverse square law. Change of orbital velocity is inversely proportional to the pressure change in the air. Pressure gradient force creates vortex gravitation.

This pattern works equally essential in a whirlwind of various volumes.

Consider the two-dimensional model from the perspective of aerodynamics. Figure 1

Figure 1: Two-dimensional model of the gravitational interaction between the two celestial bodies. The forces acting on the body 2. F_c -centrifugal force, the attractive force F_n -body 2 from the body 1, v_2 - linear velocity along the orbit of the body 2, R - Orbit radius, r_1 - radius of the body 1, r_2 - radius of the body 2, w_1 - angular velocity of rotation of ether on the surface of the body 1.



The radial distribution of pressure and velocity in the ether [4] defined on the basis of the Navier-Stokes equations for the motion of a viscous fluid (gas)

Equation 1

$$\rho \left[\frac{\partial}{\partial t} + \vec{v} \cdot \text{grad} \right] \vec{v} = \vec{F} - \text{grad } P + \eta \Delta \vec{v} \quad (1)$$

where $\rho = 8.85 \times 10^{-12} \text{ kg} \cdot \text{m}^{-3}$ - density of air - air velocity vector, P - pressure air, η - viscosity.

In cylindrical coordinates based on the radial symmetry $v_r = v_z = 0$, $v_j = v(r)$, $P = P(r)$, r -orbital radius equation written in the form of

Equation 2

$$\begin{cases} -\frac{v(r)^2}{r} = -\frac{1}{\rho} \frac{dP}{dr} \\ \eta \cdot \left(\frac{\partial^2 v(r)}{\partial r^2} + \frac{\partial v(r)}{r \partial r} - \frac{v(r)}{r^2} \right) = 0 \end{cases} \quad (2)$$

From the first equation of system (2) to be $P(r)$ for a known dependence $v(r)$, which in turn must be from the second equation. At zero viscosity, the system permits any dependence $v(r)$ [6].

Force acting on the body can be estimated by the formula

Equation 3

$$\vec{F}_\Pi = -V \cdot \text{grad } P(r) \quad (3)$$

where V - volume of the body 2.

In cylindrical coordinates for the module \vec{F}_Π

Equation 4

$$F_\Pi = V \cdot \frac{\partial P}{\partial r} \quad (4)$$

Then comparing (2) and (4) for the incompressible ether ($\rho = \text{const}$) we find that

Equation 5

$$F_\Pi = V \cdot \rho \cdot \frac{v(r)^2}{r} \quad (5)$$

Model of gravitational interaction of celestial bodies in Figure 1 can be regarded as a model of atomic attraction. In this case, the body 1 - the nucleus of an atom or an atom, which was created ethereal vortex. Each atomic vortex continuously rotates around the nucleus ester with a radius r_1 , which creates atomic force attraction. For this scheme, the gravitational forces can explain the origin of the interatomic attraction. In the theory of vortex gravitation accepted the condition that the ether permeates all matter and the body except the atomic nuclei (nucleons). Maximum speed ether several orders of magnitude greater than the speed of light. Rotation speed and the pressure gradient of ether, as well as the attractive force F_n on the surface of the atom reaches high values.

Make the assumption:

Assumption № 1: Under the influence of gravity forces on the surface atoms is compacted and / or conversion of ether into an electromagnetic flow stream including a light. Light has mass and act on it the force of gravity. Since ether, increases its density (weight), in accordance with the law of conservation of angular momentum of rotation, it should decrease in inverse proportion to its orbital speed. Thus, the electromagnetic flux received his speed - c .

When orbital revolution of the magnetic flux (light) around the atom to it are two forces: the force of the vortex atomic attraction and centrifugal force. For uniform orbital motion of light is needed to create its orbital speed centrifugal force equal to the force of an atomic attraction. This rate is

equal to the atomic orbit well-known speed of light - (c). In aerodynamics this force is called the first space.

The atomic attraction is generated by the pressure gradient in the atomic vortex (uravn.4). The pressure gradient is the difference of pressure values in free ester stationary and the center of the vortex. In a free state, the pressure is always constant and maximum. In the center of the vortex ether pressure depends on the rotational speed and air temperature. Upon heating of a substance (atom), the pressure (P) atom in the center increases in proportion to the temperature (T), in accordance with the law of Charles:

$$T \sim P \quad (6)$$

Then the pressure gradient must be reduced and, in accordance with Equation 4 reduces the force of attraction of the vortex. In this case, there comes a predominance of centrifugal force F_c over the forces of atomic, vortex attraction F_n . The speed of light is converted into the second cosmic speed. Light is removed from the atom at the most economical spiral trajectory according to calculations by Walter Hohmann. At the same time near the surface of the atom (at the lowest atomic orbits), the place flown away luminous flux takes a new ether flow. This stream is the same principle is converted into light. If the external effects (heating) is stored, the conversion and emission light from the atom will continue permanently. This is the nature of light. Light is generated and emitted not by the energy or mass of matter (atoms), but only by making the ether in the light under the influence of the atomic vortex gravitation and external influences (temperature). Light emission stop only when the external action (heating) or destruction of the atom, and with it the atomic vortex.

Figure 2 shows the movement of light aerodynamic configuration of the atom.

Figure 2: S - light source (the atom), O - observer, r - radius of the atom, r_1 - the radius of the orbit of the highest flux at his address in a calm state, prior to heating (at orbital velocity), v - relative velocity of removal or approximation between S and O, c - velocity of light vector.




Figure 2 shows: the observer (O) to move relative to the light source (S) with a speed v; light flux from the atom which trajectory intersects the observer (O) at a large angle. Consequently, the speed of light relative to the observer (O) is equal to the absolute speed of light - c. Observer speed (v) with respect to the light source can not change the rate of arrival of light (c) to the observer.

Figure 2 shows an aerodynamic scheme of movement of the light flux from a single atom. In the emission of light from the set of atoms aerodynamic design principle does not change, only multiplied many times. Light will always be supplied to each of the observer to any atom on the same helical path (at a large angle to the radius of the light field) with its constant velocity - v. Spiral light fluxes from these atoms can twist in different directions. The orbit planes of light fluxes arranged in space with any inclinations from each other. Thus, the light is almost infinite set of ultra-thin, light, spiral threads. These streams are combined into a single spherical light field. Radius and circumference of the light field in their values are directly proportional. Consequently, the light field radially increases with the same speed of light.

If the viewer moves around the light source light over a circular orbital path, and in this case a considerable part of the light will make its spiral rotation along a plane perpendicular to the plane of movement of the observer. Consequently, this light will also enter the observer at a right angle with its constant velocity - v. Isotropy of light is also a consequence of the propagation of light streams through an infinite number of spiral trajectories.

Output: To put the speed of light (c) at the speed of any body (v) in space is possible only according to the laws of vector algebra. Relative movement of any other body in space, light always moves at a right angle. Therefore, the relative velocity of the entering light to any body will always be equal to the absolute speed of light, regardless of the speed of motion of the considered point (O) relative to the light source (S).

Mathematically this can be written:

$$\bar{c} + \bar{v} = \bar{c} - \bar{v} = \bar{c}$$

CONCLUSIONS

The concept of the luminiferous ether was launched in the XVII century Rene Descartes. In the future, the wave theory, the luminiferous ether developed in the writings of Huygens. A detailed justification for this theory was in the XIX century in the framework of wave optics and electromagnetic theory of Maxwell. The proposed spiral model of light is very close to the concept of the wave motion of light and differs only in the trajectory of the light flux.

According to modern concepts light has a dual nature (wave-particle duality): - light has wave properties and is an electromagnetic wave, but is also the stream of particles - photons. It should be noted that until now scientists have not explored yet another important property of light: what is the mechanism and how can instantly disperse photons up to the maximum speed - c ?!

Spiral model of the origin and movement of light, this problem is not difficult, because the momentum of the movement all the magnetic flux received from the ether, which at the outbreak of the world itself was moving at a speed exceeding the speed of light.

Under Chapter 2 of this article can be argued that this is a consequence of the invariance of the light path of movement of the light flux. Researchers relative velocity of light in their calculations are not stacked vector velocities and their modules. The movement of the luminous flux is not contrary to the laws of classical physics. Invariance was introduced into scientific consideration only because of a misunderstanding of the movement of light as straight and radial.

REFERENCES

- [1] The relative motion of the Earth and the Luminiferous ether. Albert A. Michelson, Master, U.S.Navy The American Journal of Science. 1881. III series. Vol XXII, No. 128. P. 120—129.
- [2] Koninklijke Academie van Wetenschappen - te Amsterdam, 15, 1297 (1913)
- [3] Einstein A. «Zur Elektrodynamik bewegter Körper» Ann Phys.- 1905.- Bd 17.- S. 891
- [4] S. Orlov. Foundation of vortex gravitation, cosmology and cosmogony. Global journal of science Frontier research. Physic and Space Science Volume 12 issue 1 Version 1.0 January 2012
- [5] V A Atsurovskiy. General ether-dynamics. Energoatomizdat. Moscow, Russia. 1990. Page 278.
- [6] L V Kiknadze, Yu G Mamaladze. *Classical hydrodynamics for physicists-experimentalists*. Tbilisi University Press. Tbilisi, Georgia. 1979. Page 136.